

Gibbins, M. P., Müller, K., Glover, M., Liu, J., Putrianti, E. D., Bauza, K., Reyes-Sandoval, A., Matuschewski, K., Silvie, O. and Hafalla, J. C. R. (2020) Importance of the immunodominant CD8⁺ T cell epitope of *Plasmodium berghei* circumsporozoite protein in parasite- and vaccine-induced protection. *Infection and Immunity*, 88(10), e00383-20.

There may be differences between this version and the published version. You are advised to consult the publisher's version if you wish to cite from it.

<http://eprints.gla.ac.uk/232930/>

Deposited on: 2 February 2021

1 **The importance of the immunodominant CD8+ T cell epitope of *Plasmodium berghei***
2 **circumsporozoite protein in parasite- and vaccine-induced protection**

3

4 Matthew P. Gibbins ^a, Katja Müller ^{b,c}, Maya Glover ^a, Jasmine Liu ^a, Elyzana D. Putrianti ^{b,c},
5 Karolis Bauza ^d, Arturo Reyes-Sandoval ^d, Kai Matuschewski ^{b,c}, Olivier Silvie ^e, Julius
6 Clemence R. Hafalla ^{a#}

7

8 ^aDepartment of Infection Biology, Faculty of Infectious and Tropical Diseases, London School
9 of Hygiene and Tropical Medicine, London, United Kingdom

10 ^bParasitology Unit, Max Planck Institute for Infection Biology, Berlin, Germany

11 ^cDepartment of Molecular Parasitology, Institute of Biology, Humboldt University, Berlin,
12 Germany

13 ^dJenner Institute, Nuffield Department of Medicine, University of Oxford, United Kingdom

14 ^eSorbonne Université, INSERM, CNRS, Centre d'Immunologie et des Maladies Infectieuses,
15 CIMI-Paris, Paris, France

16

17 Running Head: Immunodominant malaria CD8+ T cell epitope

18

19 #Address correspondence to Julius Clemence R. Hafalla, Julius.Hafalla@lshtm.ac.uk

20

21 **ABSTRACT**

22 The circumsporozoite protein (CSP) builds up the surface coat of sporozoites and is the
23 leading malaria pre-erythrocytic-stage vaccine candidate. CSP has been shown to induce
24 robust CD8⁺ T cell responses that are capable of eliminating developing parasites in
25 hepatocytes resulting in protective immunity. In this study, we characterised the importance
26 of the immunodominant CSP-derived epitope, SYIPSAEKI, of *Plasmodium berghei* in both
27 sporozoite- and vaccine-induced protection in murine infection models. In BALB/c mice,
28 where SYIPSAEKI is efficiently presented in the context of the major histocompatibility
29 complex class I (MHC-I) molecule H-2-K^d, we established that epitope-specific CD8⁺ T cell
30 responses contribute to parasite killing following sporozoite immunisation. Yet, sterile
31 protection was achieved in the absence of this epitope substantiating the concept that other
32 antigens can be sufficient for parasite-induced protective immunity. Furthermore, we
33 demonstrated that SYIPSAEKI-specific CD8⁺ T cell responses elicited by viral-vectored
34 CSP-expressing vaccines effectively targeted parasites in hepatocytes. The resulting sterile
35 protection strictly relied on the expression of SYIPSAEKI. In C57BL/6 mice, which are unable
36 to present the immunodominant epitope, CSP-based vaccines did not confer complete
37 protection, despite the induction of high levels of CSP-specific antibodies. These findings
38 underscore the significance of CSP in protection against malaria pre-erythrocytic stages and
39 demonstrate that a significant proportion of the protection against the parasite is mediated by
40 CD8⁺ T cells specific for the immunodominant CSP-derived epitope.

41

INTRODUCTION

Malaria is caused by a protozoan parasite of the genus *Plasmodium* and remains a major global health challenge in tropical and subtropical countries (1). A vaccine that diminishes the burden of disease and prevents malaria transmission remains a decisive goal for malaria elimination programmes. As a gold standard in malaria vaccination, multiple immunisations of γ -radiation-attenuated *Plasmodium* sporozoites (RAS) can completely protect against wild-type (WT) sporozoite challenge (2-4). This parasite-induced protection targets the developing exo-erythrocytic forms in hepatocytes, also called liver stages, and completely abrogates blood stage infection. Antibodies and T cells have been implicated as important mechanisms of parasite-induced protection (reviewed in (5)), and CD8⁺ T cells are the prime mediators of cell-mediated protective immunity, as exemplified in murine (6, 7) and non-human primate (8) infection models.

The circumsporozoite protein (CSP), the major surface coat protein of the malaria sporozoite, has been at the forefront of vaccination studies – being the basis of RTS,S/AS01, the most progressed malaria vaccine candidate to date (9). Immunisation of BALB/c mice with *Plasmodium berghei* (*Pb*) or *P. yoelii* (*Py*) RAS evokes immunodominant major histocompatibility complex class I (MHC-I) H-2-K^d-restricted CD8⁺ T cell responses (10, 11) against distinct CSP epitopes: SYIPSAEKI for *Pb* (12) and SYVPSAEQI for *Py* (13). Indeed, the measurement of responses to these epitopes has become the standard in fundamental immunological studies in BALB/c mice and their role in parasite-induced protection from sporozoite challenge has been repeatedly demonstrated (6, 12, 14). Furthermore, numerous vaccination studies involving different viral-vectored CSP- or CSP epitope-expressing vaccines – used alone or in combination as part of prime-boost regimens – have corroborated that CSP is a highly protective antigen in the BALB/c infection model (15-21). In these studies, elevated levels of either SYIPSAEKI- or SYVPSAEQI-specific CD8⁺ T cell responses correlated with protection.

70 Nonetheless, several studies have interrogated and contested the immunological relevance
71 of CSP with regard to parasite-induced protection in contrast to vaccine-induced protection.
72 These studies emanated from observations that in naturally exposed humans T cell
73 responses to CSP are scarce (22). In murine malaria models, multiple immunisations are
74 required to elicit CD8+ T cell-dependent protective immunity in various mouse strains,
75 particularly where no other strongly immunogenic CSP-derived CD8+ T cell epitopes have
76 been identified (23). Furthermore, in *Py*CSP-transgenic BALB/c mice that are tolerant to
77 *Py*CSP, complete protection can be achieved by *Py* RAS immunisation (24). In good
78 agreement, BALB/c mice immunised with *Pb* WT parasites are completely protected when
79 challenged with transgenic *Pb* parasites where the endogenous CSP has been swapped with
80 the *P. falciparum* CSP (25). Taken together, these studies indicate that immune responses to
81 CSP are dispensable for protection, and that other antigens are important to elicit protective
82 immunity.

83

84 In this study, we have extended previous work on the entire CSP by dissecting the relevance
85 of a single CSP-derived immunodominant epitope in parasite- and vaccine-induced
86 protection. As the most stringent model system, we utilised transgenic *Pb* parasites lacking
87 SYIPSAEKI for immunisation and challenge experiments in BALB/c mice. In addition, we
88 have highlighted the level of protection achieved by CSP-based vaccines in mice expressing
89 the relevant (BALB/c) or irrelevant (C57BL/6) MHC-I needed to present the CSP-derived
90 immunodominant epitope.

91

92 RESULTS

93 Sporozoite-induced SYIPSAEKI-specific CD8⁺ T cell responses contribute to parasite 94 killing but are dispensable for the development of sterile immunity.

95 First, we interrogated the role that SYIPSAEKI, the H-2-K^d-restricted immunodominant
96 epitope of *PbCSP*, plays in protective immunity induced after live attenuated sporozoite
97 immunisation. For this purpose, *PbCSP*^{SIINFEKL} radiation-attenuated sporozoites (RAS), where
98 the SYPSAEKI sequence has been replaced with the H-2-K^b-restricted epitope of ovalbumin,
99 SIINFEKL ((26), Müller and Gibbins et al., unpublished) were used to immunise H-2-K^d-
100 expressing BALB/c mice. To date, there are no other reported strongly immunogenic H-2-K^d-
101 restricted *PbCSP* epitopes identified. Removal of SYPSAEKI, by replacement with an
102 irrelevant epitope, in the *PbCSP*^{SIINFEKL} parasites allows unequivocal assignment of critical
103 roles of this immunodominant *PbCSP*-derived epitope in protection elicited by live sporozoite
104 immunisations. Two weeks after immunisation, the frequencies of IFN- γ -producing
105 SYIPSAEKI-specific CD8⁺ T cell responses in the spleen after gating for CD11a expression
106 (an activation marker commonly used to identify antigen-experienced cells (27)) were
107 measured by flow cytometry (Fig. 1A). As expected, *PbCSP*^{SIINFEKL} RAS parasites elicited no
108 SYIPSAEKI-specific CD8⁺ T cell responses in BALB/c mice.

109

110 To ascertain whether SYIPSAEKI contributes to parasite-induced protection, BALB/c mice
111 were immunised once (Fig 1B) or twice (Fig 1C) with either *PbWT* or *PbCSP*^{SIINFEKL} RAS.
112 Two weeks after immunisation, the mice were challenged with *PbWT* sporozoites and
113 protection was determined by measuring the parasite loads in the liver 40 hours later (Fig. 1B
114 and C). A significant reduction in parasite load – up to four orders of magnitude difference as
115 compared to naïve mice – was observed in mice immunised with *PbWT* RAS and challenged
116 with *PbWT* parasites. In contrast, protection was reduced only by approximately two orders
117 of magnitude (after one immunisation) or three orders of magnitude (after two immunisations)
118 in mice immunised with *PbCSP*^{SIINFEKL} RAS (Fig. 1B and C). These results highlight the
119 notion that within *PbCSP*, the SYIPSAEKI epitope has a critical and immunodominant
120 contribution to protecting BALB/c mice after one or two immunisations with RAS.

121

122 However, multiple immunisations with RAS are required to induce sterile protection. To
123 establish whether the development of sterile immunity is dependent on SYIPSAEKI-specific
124 CD8⁺ T cell responses, BALB/c mice were immunised thrice with *PbCSP*^{SIINFEKL} RAS one
125 week apart; two weeks after the last immunisation, mice were challenged with *PbWT*
126 sporozoites (Fig. 1C). All mice were protected from blood stage infection compared to the
127 naïve controls, implying that SYIPSAEKI-specific CD8⁺ T cell responses are not necessary
128 for the development of sterile immunity.

129

130 **Prime-boost vaccination with CSP-expressing viruses induces strong anti-CSP**
131 **antibody and CD8⁺ T cell responses and SYIPSAEKI is the key mediator of sterile**
132 **protection.**

133 Next, we probed the requirement for SYIPSAEKI presentation in protection elicited by viral-
134 vectored CSP-expressing vaccines administered in a prime-boost regimen. Priming with
135 adenovirus (Ad) carrying a foreign antigen and boosting with orthopoxvirus modified vaccinia
136 Ankara (MVA) expressing the same antigen has consistently been shown to induce strong
137 CD8⁺ T cell responses with high levels of protective efficacy against intracellular pathogens
138 including malaria pre-erythrocytic stages (18, 21).

139

140 Chimpanzee adenovirus serotype 63 (AdCh63) and MVA vaccines expressing *PbCSP* were
141 used to vaccinate BALB/c mice with a two-week resting period between priming and boosting
142 (Fig. 2A). Two weeks after boosting, whole blood was collected and restimulated *ex vivo* with
143 SYIPSAEKI peptide. The frequencies of IFN- γ secreting CD8⁺ T cells were enumerated by
144 flow cytometry (Fig. 2B) and Ad-MVA *PbCSP*-vaccinated mice elicited ~12% SYIPSAEKI-
145 specific circulating CD8⁺ T cells (Fig. 2C), which is consistent with the level of epitope-
146 specific CD8⁺ T cells induced by similar, previously published, viral vectored prime-boost
147 regimens (19, 21). Serum samples were also collected from the vaccinated animals and
148 were used in an immunofluorescence assay against air-dried *Pb* sporozoites (Fig. 2D). Ad-
149 MVA *PbCSP*-vaccinated BALB/c mice induced high anti-CSP antibody titres (1:10⁴). These

150 data indicate that Ad-MVA *PbCSP* vaccination elicit both high frequencies of SYPSAEKI-
151 specific CD8+ T cells and high titres of CSP-specific antibodies.

152

153 Two weeks after boosting, Ad-MVA *PbCSP*-vaccinated mice were challenged with *PbWT* or
154 *PbCSP*^{SIINFEKL} parasites. Protection was assessed by two complementary assays; (i)
155 determination of the reduction of parasite load in the liver (Fig. 2E) and (ii) induction of sterile
156 protection (Fig. 2F). Strikingly, parasite load in the liver of Ad-MVA *PbCSP*-vaccinated mice
157 was not significantly reduced compared to non-vaccinated mice when challenged with
158 *PbCSP*^{SIINFEKL} sporozoites, in marked contrast to challenge with *PbWT* sporozoites. In perfect
159 agreement, vaccinated mice challenged with *PbCSP*^{SIINFEKL} sporozoites were patent for
160 parasitaemia by day 5, whereas vaccinated mice challenged with *PbWT* sporozoites
161 remained completely protected. These results denote that vaccine-induced effector
162 SYIPSEAKI-specific CD8+ T responses efficiently target parasites expressing the cognate
163 epitope. Parasites lacking the SYIPSEAKI epitope are not eliminated despite high levels of
164 CSP-specific antibodies evoked by vaccination in this experimental system.

165

166 **CSP-based vaccines do not elicit protective immunity in C57BL/6 mice.**

167 To further investigate the requirement of SYIPSEAKI as the indispensable protective epitope
168 of CSP, mice unable to present this epitope were vaccinated with the *PbCSP* prime-boost
169 regimen with an interval of two weeks between vaccines, followed by challenge with either
170 *PbWT* or *PbCSP*^{SIINFEKL} parasites (Fig 3A). C57BL/6 mice were used because SYIPSEAKI is
171 an H-2-K^d restricted epitope, and this mouse strain does not express the relevant MHC-I
172 allele. Thus, SYIPSEAKI would fail to be presented by infected hepatocytes. As before, blood
173 and serum were derived two weeks after boost. As expected, SYIPSEAKI-specific CD8+ T
174 cells (Fig. 3B) were not detectable in Ad-MVA *PbCSP*-vaccinated C57BL/6 mice (Fig. 3C),
175 but strong anti-CSP antibody titres (1:10⁴) were elicited (Fig. 3D). Ad-MVA CSP-vaccinated
176 C57BL/6 mice challenged with either *PbWT* or *PbCSP*^{SIINFEKL} parasites had comparable
177 parasite load in the liver (Fig. 3E), indicative of full liver stage development in all groups.

178

179 **DISCUSSION**

180 Our findings lend full support to the notion that CSP is an immunodominant sporozoite-
181 derived antigen (24). A single epitope, SYIPSAEKI, is the immunodominant CD8+ T cell
182 epitope of CSP, and we show that it is responsible for the antigen's protective capacity
183 against parasites in the liver in the BALB/c model. Following RAS immunisation, CD8+ T cell
184 responses to SYIPSAEKI contribute to the reduction in parasite load in the liver following
185 sporozoite challenge, as shown herein. When RAS-immunised mice are challenged with
186 *PbCSP*^{SIINFEKL}, transgenic parasites lacking SYIPSAEKI, reduced anti-*Plasmodium* activity in
187 the liver is observed. Nonetheless, complete protection is achievable in the absence of
188 SYIPSAEKI-specific CD8+ T cell responses, demonstrating that responses to other, yet
189 unidentified, H-2-K^d-restricted epitopes contribute to parasite killing. It is conceivable that
190 these epitopes are encoded by the hundreds of other *Plasmodium* genes expressed in
191 malaria pre-erythrocytic stages, some of which might be shared with blood stage antigens
192 (28).

193
194 Our findings also emphasise the importance of SYIPSAEKI-specific CD8+ T cell responses
195 for promoting protective immunity when using CSP-based viral vaccines in the BALB/c
196 model. These vaccines are aimed at generating high levels of epitope-specific memory CD8+
197 T cells but rely on the expression of relevant MHC-I in the vaccinated host and the presence
198 of the cognate epitope in the parasite used for challenge (29). Notably, despite high levels of
199 antibodies against whole sporozoites elicited following Ad-MVA *PbCSP* vaccination, sterile
200 protection was not achieved following challenge of C57BL/6 mice. These mice cannot
201 present SYIPSAEKI, fully supporting the notion that the protective efficacy of CSP strictly
202 depends on the expression of the immunodominant epitope. These findings were
203 independently corroborated by the lack of protection in mice, either BALB/c or C57BL/6,
204 immunised with transgenic sporozoites lacking SYIPSAEKI.

205

206 Together, these results have important implications for the development of next generation
207 malaria vaccines. We have demonstrated the significance of a single epitope of CSP in

208 mediating protective CD8⁺ T cell responses while also recapitulating that protection can be
209 achieved in the absence of responses to the entire CSP antigen (24, 25). In BALB/c mice,
210 SYIPSAEKI-specific CD8⁺ T cell responses offered protection. However, to achieve
211 complete sterile protection either multiple sporozoite immunisations or viral vaccines, which
212 induced large populations of SYIPSAEKI-specific CD8⁺ T cells, were required. Multiple
213 immunisations likely induced a broad range of immune responses and multiple high-dose
214 immunisations with RAS in humans have been shown to induce dose-dependent anti-
215 sporozoite CD8⁺ T cell responses in addition to dose dependent anti-sporozoite antibody
216 and CD4⁺ T cell responses (4). It will be important in the future to determine how the
217 magnitude of SYIPSAEKI-specific CD8⁺ T cell responses modulates, after consecutive
218 immunisations, but also the breadth of responses to other CSP epitopes (B and T cell) and
219 their effect on protection compared to other antigens. In line with this, our findings also show
220 that protection can be achieved in the absence of responses to immunodominant epitopes,
221 leading us to suggest that future pre-erythrocytic malaria vaccine research should not only
222 focus on inducing strong CD8⁺ T cell responses against one or multiple antigens but should
223 try to target a broad array of antigens and cover diverse MHC to offer the best protection
224 possible. The identification of novel antigens and epitopes that contribute to protection in H-
225 2-K^d-restricted BALB/c mice, and ultimately in human populations with broad MHC
226 haplotypes, will aid this development. In C57BL/6 mice pre-erythrocytic immunity is mounted
227 irrespective of CSP-specific CD8⁺ T cell responses, and recent genome-wide epitope
228 profiling returned multiple sporozoite antigens and epitopes (30-32). RTS,S/AS01, the
229 leading subunit malaria vaccine based on CSP, seems to offer some protection against *P.*
230 *falciparum* re-infection (9). Partial and short-lived protection is likely primarily mediated by the
231 action of transitory anti-sporozoite antibodies (33-35). Strikingly, peripheral blood CD8⁺ T cell
232 responses were not identified to provide a role following sporozoite challenge in this
233 candidate vaccine. Together with previous findings (7, 17, 19, 24) our data underscore efforts
234 to improve the most advanced candidate malaria vaccine, RTS,S/AS01, by eliciting CD8⁺ T
235 cells against CSP or other immunodominant antigens.

236

237 **MATERIALS AND METHODS**

238 **Ethics and animal experimentation.** Animal procedures were performed in accordance
239 with the German 'Tierschutzgesetz in der Fassung vom 18. Mai 2006 (BGBl. I S. 1207)'
240 which implements the directive 2010/6 3/EU from the European Union. The protocol was
241 approved by the ethics committee of the Berlin state authority ('Landesamt für Gesundheit
242 und Soziales Berlin', permit number G0469/09). Animal experiments at London School of
243 Hygiene and Tropical Medicine were conducted under license from the United Kingdom
244 Home Office under the Animals (Scientific Procedures) Act 1986. CD-1 mice were bred in-
245 house at LSHTM, while NMRI, C57BL/6 and BALB/c laboratory mouse strains were
246 purchased from either Charles River Laboratories (Margate, UK or Sulzfeld, Germany) or
247 Janvier (Saint Berthevin, France). Female mice of 6-8 weeks of age were used in the
248 experiments.

249

250 **Plasmodium parasites and immunisation.** The transgenic *P. berghei* ANKA CSP^{SIINFEKL}
251 (*PbCSP*^{SIINFEKL}) parasite was generated with the immunodominant CSP CD8+ T cell epitope
252 SYIPSAEKI (252-260aa) being replaced with the H-2^b restricted *Gallus gallus* ovalbumin
253 CD8+ T cell epitope SIINFEKL (258-265aa) via double homologous recombination ((26),
254 Müller and Gibbins *et al.*, unpublished). Wild-type *Plasmodium berghei* ANKA (clone
255 c115cy1) (*PbWT*) and *PbCSP*^{SIINFEKL} were maintained by continuous cycling between murine
256 hosts (NMRI or CD-1) and *Anopheles stephensi* mosquitos. Infected mosquitos were kept in
257 incubators (Panasonic and Mytron) at 80% humidity and 20°C temperature. Sporozoites
258 were isolated from the salivary glands and attenuated by γ -irradiation at 1.2x10⁴cGy. Mice
259 were immunised with 10,000 sporozoites administered intravenously with multiple doses
260 given one week apart unless otherwise stated. For challenge infections, 5,000 or 10,000
261 sporozoites were administered intravenously to assess sterile protection and parasite load in
262 the liver, respectively.

263

264 **Viral-vectored CSP-expressing vaccines.** AdCh63 and MVA vaccines expressing the
265 mammalian codon-optimised fragment of *PbCSP* were constructed and propagated based

on previously published viral vectors (36, 37). The viral vectors were administered intramuscularly in endotoxin-free PBS at a concentration of 10^5 viral particles for AdPbCSP for the prime immunisation and 10^6 viral particles MVAPbCSP for the boost immunisation.

Immunofluorescent antibody assay. 10,000 sporozoites were spotted onto epoxy coated glass slides with marked rings (Medco), dried at room temperature and stored at -20°C . Thawed slides were fixed in acetone, dried and rehydrated with PBS before incubation in 10% FCS supplemented DMEM (Gibco) for 1 hour at 37°C in a humid chamber. Serum at concentrations $1:10^3$, $1:3.3 \times 10^3$, $1:10^4$, $1:3.3 \times 10^4$, $1:10^5$ (and, additionally, $1:3.3 \times 10^5$ and $1:10^6$ for C57BL/6 serum) were added to the ring wells and incubated for 1 hour at 37°C in a humid chamber. Slides were washed and stained with a mouse anti-CSP (38) primary antibody. Hoechst33342 was added as the nuclear stain together with a respective fluorescently labelled anti-mouse secondary antibody for a further one-hour incubation. Slides were washed and mounted with 'Fluoromount-G' (Southern Biotech) and analysed by fluorescent microscopy (Zeiss Axio Observer).

Quantification of SYIPSAEKI-specific CD8+ T cell responses. Spleens were harvested and lymphocytes were derived by passing spleens through $40\mu\text{m}$ cell strainers (Corning). Peripheral blood was drawn from the tail vein and collected in Na^+ heparin capillary tubes (Brand) and assayed in 96-well flat bottom plates (Corning). Red blood cells were lysed using PharmLyse (BD) and lymphocytes were resuspended in 10% FCS, 2% Penicillin-Streptomycin and 1% L-glutamine supplemented RPMI 1640 (Gibco). Splenocytes were counted using a 40x dilution with Trypan Blue (ThermoFisher Scientific) and a Neubauer 'Improved' haemocytometer (Biochrom). 2×10^6 splenocytes and the lysed blood samples were prepared in 96 well plates and incubated with a final concentration of $10\mu\text{g/ml}$ of SYIPSAEKI peptide in the presence of Brefeldin A (eBioScience) for 5-6 hours at 37°C and 5% CO_2 . For staining of cell surface markers and intracellular cytokines, cells were incubated for 1 hour at 4°C for each staining. Cells were stained for CD8 (53-6.7) and CD11a

294 (M17/4) (eBioscience). Splenic cells were fixed with 4% paraformaldehyde and peripheral
295 blood cells were fixed with 1% paraformaldehyde before staining for IFN- γ (XMG1.2)
296 (eBioscience) in the presence of Perm/Wash buffer (BD) for intracellular staining. Data was
297 acquired by flow cytometry using an LSRFortessa or LSRII (BD) and analysed using
298 Flowjo9.5.2 (Tree Star, Inc.).

299

300 **Quantification of parasite load in the liver.** Livers were harvested 40-42 hours after
301 sporozoite challenge and total RNA was extracted following homogenisation using TRIzol
302 (ThermoFisher Scientific). cDNA was generated using the RETROScript Kit (Ambion).
303 Quantitative real-time PCR was performed using the StepOnePlus Real-Time PCR System
304 and Power SYBR Green PCR Master Mix (Applied Biosystems). Relative liver parasite levels
305 were quantified using the $\Delta\Delta C_t$ method comparing levels of *P. berghei* 18S rRNA using
306 specific primers and normalised to levels of mouse *GAPDH* mRNA (39).

307

308 **Assessment of parasitaemia.** Sterile protection was assessed by daily blood smears, taken
309 from mice 3-14 days after sporozoite challenge, stained with Giemsa (improved solution;
310 VWR) to microscopically determine the presence of blood stage parasites.

311

312 **Statistical analysis.** Statistical analysis was performed using GraphPad Prism v7
313 (GraphPad Software, Inc.). Statistics were calculated using the Mann-Whitney U test.

314

315 **AUTHOR CONTRIBUTIONS**

316 O.S. and J.C.R.H. designed the experiments in the laboratory of K.Matuschewski; O.S.
317 generated the transgenic parasites CSP^{SIINFEKL}; M.P.G., K.Müller., M.G., J.L. and E.D.P.
318 performed experiments and analysed data; K.B. and A.R.-S. generated the CSP-expressing
319 viruses AdPbCSP and MVAPbCSP; M.P.G. and J.C.R.H. wrote the paper. All authors
320 commented on and approved the paper.

321

322 **ACKNOWLEDGEMENTS**

323 J.C.R.H. was funded by grants from The Royal Society (University Research Fellowship
324 UF0762736/UF120026 and Project Grant RG130034) and the National Centre for the
325 Replacement, Refinement & Reduction of Animals in Research (Project Grant
326 NC/L000601/1). O.S. was funded in part by the Laboratoire d'Excellence ParaFrap (ANR-11-
327 LABX-0024). K.Matuschewski was supported by the Max Planck Society and grants from the
328 European Commission (EviMalaR Network of Excellence #34), the Chica and Heinz Schaller
329 Foundation, and the Alliance Berlin Canberra "Crossing Boundaries: Molecular Interactions
330 in Malaria", which is co-funded by a grant from the Deutsche Forschungsgemeinschaft (DFG)
331 for the International Research Training Group (IRTG) 2290 and the Australian National
332 University. A.R.-S., a Jenner Investigator and an Oxford Martin Fellow, was funded by
333 a Wellcome Trust Career Development Fellowship (Grant 097395/Z/11/Z). We would like to
334 thank the Jenner Institute's Viral Vector Core Facility for the viral vectored vaccines. The
335 funders had no role in study design, data collection and analysis, decision to publish, or
336 preparation of the manuscript.

337 REFERENCES

338

- 339 1. WHO. 2017. World Malaria Report 2017 World Health Organisation, Geneva, Switzerland.
- 340 2. Nussenzweig RS, Vanderberg J, Most H, Orton C. 1967. Protective immunity produced by the
341 injection of x-irradiated sporozoites of *Plasmodium berghei*. *Nature* 216:160-2.
- 342 3. Hoffman SL, Goh LM, Luke TC, Schneider I, Le TP, Doolan DL, Sacchi J, de la Vega P, Dowler M,
343 Paul C, Gordon DM, Stoute JA, Church LW, Sedegah M, Heppner DG, Ballou WR, Richie TL.
344 2002. Protection of humans against malaria by immunization with radiation-attenuated
345 *Plasmodium falciparum* sporozoites. *J Infect Dis* 185:1155-64.
- 346 4. Seder RA, Chang LJ, Enama ME, Zephir KL, Sarwar UN, Gordon IJ, Holman LA, James ER,
347 Billingsley PF, Gunasekera A, Richman A, Chakravarty S, Manoj A, Velmurugan S, Li M, Ruben
348 AJ, Li T, Eappen AG, Stafford RE, Plummer SH, Hendel CS, Novik L, Costner PJ, Mendoza FH,
349 Saunders JG, Nason MC, Richardson JH, Murphy J, Davidson SA, Richie TL, Sedegah M,
350 Sutamihardja A, Fahle GA, Lyke KE, Laurens MB, Roederer M, Tewari K, Epstein JE, Sim BK,
351 Ledgerwood JE, Graham BS, Hoffman SL, Team VRCS. 2013. Protection against malaria by
352 intravenous immunization with a nonreplicating sporozoite vaccine. *Science* 341:1359-65.
- 353 5. Doll KL, Harty JT. 2014. Correlates of protective immunity following whole sporozoite
354 vaccination against malaria. *Immunol Res* 59:166-76.
- 355 6. Doolan DL, Hoffman SL. 2000. The complexity of protective immunity against liver-stage
356 malaria. *J Immunol* 165:1453-62.
- 357 7. Schmidt NW, Podyminogin RL, Butler NS, Badovinac VP, Tucker BJ, Bahjat KS, Lauer P, Reyes-
358 Sandoval A, Hutchings CL, Moore AC, Gilbert SC, Hill AV, Bartholomay LC, Harty JT. 2008.
359 Memory CD8 T cell responses exceeding a large but definable threshold provide long-term
360 immunity to malaria. *Proc Natl Acad Sci U S A* 105:14017-22.
- 361 8. Weiss WR, Jiang CG. 2012. Protective CD8+ T lymphocytes in primates immunized with
362 malaria sporozoites. *PLoS One* 7:e31247.
- 363 9. RTS SCTP. 2015. Efficacy and safety of RTS,S/AS01 malaria vaccine with or without a booster
364 dose in infants and children in Africa: final results of a phase 3, individually randomised,
365 controlled trial. *Lancet* 386:31-45.
- 366 10. Schofield L, Villaquiran J, Ferreira A, Schellekens H, Nussenzweig R, Nussenzweig V. 1987.
367 Gamma interferon, CD8+ T cells and antibodies required for immunity to malaria sporozoites.
368 *Nature* 330:664-6.
- 369 11. Weiss WR, Sedegah M, Beaudoin RL, Miller LH, Good MF. 1988. CD8+ T cells
370 (cytotoxic/suppressors) are required for protection in mice immunized with malaria
371 sporozoites. *Proc Natl Acad Sci U S A* 85:573-6.
- 372 12. Romero P, Maryanski JL, Corradin G, Nussenzweig RS, Nussenzweig V, Zavala F. 1989. Cloned
373 cytotoxic T cells recognize an epitope in the circumsporozoite protein and protect against
374 malaria. *Nature* 341:323-6.
- 375 13. Weiss WR, Mellouk S, Houghten RA, Sedegah M, Kumar S, Good MF, Berzofsky JA, Miller LH,
376 Hoffman SL. 1990. Cytotoxic T cells recognize a peptide from the circumsporozoite protein on
377 malaria-infected hepatocytes. *J Exp Med* 171:763-73.
- 378 14. Rodrigues MM, Cordey AS, Arreaza G, Corradin G, Romero P, Maryanski JL, Nussenzweig RS,
379 Zavala F. 1991. CD8+ cytolytic T cell clones derived against the *Plasmodium yoelii*
380 circumsporozoite protein protect against malaria. *Int Immunol* 3:579-85.
- 381 15. Li S, Rodrigues M, Rodriguez D, Rodriguez JR, Esteban M, Palese P, Nussenzweig RS, Zavala F.
382 1993. Priming with recombinant influenza virus followed by administration of recombinant
383 vaccinia virus induces CD8+ T-cell-mediated protective immunity against malaria. *Proc Natl*
384 *Acad Sci U S A* 90:5214-8.
- 385 16. Rodrigues EG, Zavala F, Eichinger D, Wilson JM, Tsuji M. 1997. Single immunizing dose of
386 recombinant adenovirus efficiently induces CD8+ T cell-mediated protective immunity
387 against malaria. *J Immunol* 158:1268-74.

- 388 17. Bruna-Romero O, Gonzalez-Aseguinolaza G, Hafalla JC, Tsuji M, Nussenzweig RS. 2001.
389 Complete, long-lasting protection against malaria of mice primed and boosted with two
390 distinct viral vectors expressing the same plasmodial antigen. *Proc Natl Acad Sci U S A*
391 98:11491-6.
- 392 18. Gilbert SC, Schneider J, Hannan CM, Hu JT, Plebanski M, Sinden R, Hill AV. 2002. Enhanced
393 CD8 T cell immunogenicity and protective efficacy in a mouse malaria model using a
394 recombinant adenoviral vaccine in heterologous prime-boost immunisation regimes. *Vaccine*
395 20:1039-45.
- 396 19. Gonzalez-Aseguinolaza G, Nakaya Y, Molano A, Dy E, Esteban M, Rodriguez D, Rodriguez JR,
397 Palese P, Garcia-Sastre A, Nussenzweig RS. 2003. Induction of protective immunity against
398 malaria by priming-boosting immunization with recombinant cold-adapted influenza and
399 modified vaccinia Ankara viruses expressing a CD8+ T-cell epitope derived from the
400 circumsporozoite protein of *Plasmodium yoelii*. *J Virol* 77:11859-66.
- 401 20. Reyes-Sandoval A, Sridhar S, Berthoud T, Moore AC, Harty JT, Gilbert SC, Gao G, Ertl HC,
402 Wilson JC, Hill AV. 2008. Single-dose immunogenicity and protective efficacy of simian
403 adenoviral vectors against *Plasmodium berghei*. *Eur J Immunol* 38:732-41.
- 404 21. Reyes-Sandoval A, Berthoud T, Alder N, Siani L, Gilbert SC, Nicosia A, Colloca S, Cortese R, Hill
405 AV. 2010. Prime-boost immunization with adenoviral and modified vaccinia virus Ankara
406 vectors enhances the durability and polyfunctionality of protective malaria CD8+ T-cell
407 responses. *Infect Immun* 78:145-53.
- 408 22. Dodoo D, Hollingdale MR, Anum D, Koram KA, Gyan B, Akanmori BD, Ocran J, Adu-
409 Amankwah S, Geneshan H, Abot E, Legano J, Banania G, Sayo R, Brambilla D, Kumar S, Doolan
410 DL, Rogers WO, Epstein J, Richie TL, Sedegah M. 2011. Measuring naturally acquired immune
411 responses to candidate malaria vaccine antigens in Ghanaian adults. *Malar J* 10:168.
- 412 23. Schmidt NW, Butler NS, Badovinac VP, Harty JT. 2010. Extreme CD8 T cell requirements for
413 anti-malarial liver-stage immunity following immunization with radiation attenuated
414 sporozoites. *PLoS Pathog* 6:e1000998.
- 415 24. Kumar KA, Sano G, Boscardin S, Nussenzweig RS, Nussenzweig MC, Zavala F, Nussenzweig V.
416 2006. The circumsporozoite protein is an immunodominant protective antigen in irradiated
417 sporozoites. *Nature* 444:937-40.
- 418 25. Gruner AC, Mauduit M, Tewari R, Romero JF, Depinay N, Kayibanda M, Lallemand E, Chavatte
419 JM, Crisanti A, Sinnis P, Mazier D, Corradin G, Snounou G, Renia L. 2007. Sterile protection
420 against malaria is independent of immune responses to the circumsporozoite protein. *PLoS*
421 *One* 2:e1371.
- 422 26. Cockburn IA, Tse SW, Radtke AJ, Srinivasan P, Chen YC, Sinnis P, Zavala F. 2011. Dendritic cells
423 and hepatocytes use distinct pathways to process protective antigen from plasmodium in
424 vivo. *PLoS Pathog* 7:e1001318.
- 425 27. Rai D, Pham NL, Harty JT, Badovinac VP. 2009. Tracking the total CD8 T cell response to
426 infection reveals substantial discordance in magnitude and kinetics between inbred and
427 outbred hosts. *J Immunol* 183:7672-81.
- 428 28. Muller K, Gibbins MP, Matuschewski K, Hafalla JCR. 2017. Evidence of cross-stage CD8+ T cell
429 epitopes in malaria pre-erythrocytic and blood stage infections. *Parasite Immunol* 39.
- 430 29. Silvie O, Amino R, Hafalla JC. 2017. Tissue-specific cellular immune responses to malaria pre-
431 erythrocytic stages. *Curr Opin Microbiol* 40:160-167.
- 432 30. Hafalla JC, Bauza K, Friesen J, Gonzalez-Aseguinolaza G, Hill AV, Matuschewski K. 2013.
433 Identification of targets of CD8(+) T cell responses to malaria liver stages by genome-wide
434 epitope profiling. *PLoS Pathog* 9:e1003303.
- 435 31. Lau LS, Fernandez-Ruiz D, Mollard V, Sturm A, Neller MA, Cozijnsen A, Gregory JL, Davey GM,
436 Jones CM, Lin YH, Haque A, Engwerda CR, Nie CQ, Hansen DS, Murphy KM, Papenfuss AT,
437 Miles JJ, Burrows SR, de Koning-Ward T, McFadden GI, Carbone FR, Crabb BS, Heath WR.
438 2014. CD8+ T cells from a novel T cell receptor transgenic mouse induce liver-stage immunity
439 that can be boosted by blood-stage infection in rodent malaria. *PLoS Pathog* 10:e1004135.

- 440 32. Pichugin A, Zarling S, Perazzo L, Duffy PE, Ploegh HL, Krzych U. 2018. Identification of a Novel
441 CD8 T Cell Epitope Derived from *Plasmodium berghei* Protective Liver-Stage Antigen. *Front*
442 *Immunol* 9:91.
- 443 33. Olotu A, Lusingu J, Leach A, Lievens M, Vekemans J, Msham S, Lang T, Gould J, Dubois MC,
444 Jongert E, Vansadia P, Carter T, Njuguna P, Awuondo KO, Malabeja A, Abdul O, Gesase S,
445 Mturi N, Drakeley CJ, Savarese B, Villafana T, Lapierre D, Ballou WR, Cohen J, Lemnge MM,
446 Peshu N, Marsh K, Riley EM, von Seidlein L, Bejon P. 2011. Efficacy of RTS,S/AS01E malaria
447 vaccine and exploratory analysis on anti-circumsporozoite antibody titres and protection in
448 children aged 5-17 months in Kenya and Tanzania: a randomised controlled trial. *Lancet*
449 *Infect Dis* 11:102-9.
- 450 34. White MT, Verity R, Griffin JT, Asante KP, Owusu-Agyei S, Greenwood B, Drakeley C, Gesase
451 S, Lusingu J, Ansong D, Adjei S, Agbenyega T, Ogutu B, Otieno L, Otieno W, Agnandji ST, Lell B,
452 Kremsner P, Hoffman I, Martinson F, Kamthunzu P, Tinto H, Valea I, Sorgho H, Oneko M,
453 Otieno K, Hamel MJ, Salim N, Mtoro A, Abdulla S, Aide P, Sacarlal J, Aponte JJ, Njuguna P,
454 Marsh K, Bejon P, Riley EM, Ghani AC. 2015. Immunogenicity of the RTS,S/AS01 malaria
455 vaccine and implications for duration of vaccine efficacy: secondary analysis of data from a
456 phase 3 randomised controlled trial. *Lancet Infect Dis* 15:1450-8.
- 457 35. Kester KE, Cummings JF, Ofori-Anyinam O, Ockenhouse CF, Krzych U, Moris P, Schwenk R,
458 Nielsen RA, Debebe Z, Pinelis E, Juompan L, Williams J, Dowler M, Stewart VA, Wirtz RA,
459 Dubois MC, Lievens M, Cohen J, Ballou WR, Heppner DG, Jr., Rts SVEG. 2009. Randomized,
460 double-blind, phase 2a trial of falciparum malaria vaccines RTS,S/AS01B and RTS,S/AS02A in
461 malaria-naïve adults: safety, efficacy, and immunologic associates of protection. *J Infect Dis*
462 200:337-46.
- 463 36. O'Hara GA, Duncan CJ, Ewer KJ, Collins KA, Elias SC, Halstead FD, Goodman AL, Edwards NJ,
464 Reyes-Sandoval A, Bird P, Rowland R, Sheehy SH, Poulton ID, Hutchings C, Todryk S, Andrews
465 L, Folgori A, Berrie E, Moyle S, Nicosia A, Colloca S, Cortese R, Siani L, Lawrie AM, Gilbert SC,
466 Hill AV. 2012. Clinical assessment of a recombinant simian adenovirus ChAd63: a potent new
467 vaccine vector. *J Infect Dis* 205:772-81.
- 468 37. Reyes-Sandoval A, Wyllie DH, Bauza K, Milicic A, Forbes EK, Rollier CS, Hill AV. 2011. CD8+ T
469 effector memory cells protect against liver-stage malaria. *J Immunol* 187:1347-57.
- 470 38. Potocnjak P, Yoshida N, Nussenzweig RS, Nussenzweig V. 1980. Monovalent fragments (Fab)
471 of monoclonal antibodies to a sporozoite surface antigen (Pb44) protect mice against
472 malarial infection. *J Exp Med* 151:1504-13.
- 473 39. Muller K, Matuschewski K, Silvie O. 2011. The Puf-family RNA-binding protein Puf2 controls
474 sporozoite conversion to liver stages in the malaria parasite. *PLoS One* 6:e19860.
475
476

477 **FIGURE LEGENDS**

478 **FIG 1** SYIPSAEKI is dispensable for RAS immunisation but predominates protection with
479 fewer immunisations. (A) BALB/c mice were immunised once with 10,000 *PbWT* or
480 *PbCSP*^{SIINFEKL} RAS. Splenocytes were taken after two weeks and restimulated with
481 SYIPSAEKI peptide. IFN- γ -producing cells co-staining with CD11a were assessed by flow
482 cytometry. Shown are the time course (top), the gating strategy (centre) and proportion of
483 IFN- γ -producing CD11a of total CD8+ T cells (bottom). (B) Groups of BALB/c mice were
484 immunised once with 15,000 *PbWT* or *PbCSP*^{SIINFEKL} RAS. Immunised mice and BALB/c
485 naïve controls (n=3-10) were challenged with 10,000 *PbWT* parasites two weeks after the
486 last immunisation. Livers were harvested 40 hours post-challenge and the relative liver
487 parasite loads were quantified using the $\Delta\Delta$ Ct method comparing levels of *P. berghei* 18S
488 rRNA and levels of mouse *GAPDH* mRNA. Mean values (\pm SEM) are shown and statistics
489 were calculated using the Kruskal-Wallis test (***, p<0.001). (C) Groups of BALB/c mice
490 were immunised twice with 10,000 *PbWT* or *PbCSP*^{SIINFEKL} RAS, 1 week apart. Immunised
491 mice and BALB/c naïve controls (n=5) were challenged with 10,000 *PbWT* parasites two
492 weeks after the last immunisation. Livers were harvested 40 hours post-challenge and the
493 relative liver parasite loads were quantified using the $\Delta\Delta$ Ct method comparing levels of *P.*
494 *berghei* 18S rRNA and levels of mouse *GAPDH* mRNA. Mean values (\pm SEM) are shown and
495 statistics were calculated using the Kruskal-Wallis test (***, p<0.001). (D) BALB/c mice
496 (n=12) were immunised with three doses of 10,000 *PbCSP*^{SIINFEKL} RAS at one-week intervals.
497 Immunised mice and naïve controls (n=11) were challenged with 5,000 *PbWT* sporozoites 16
498 days after the last immunisation. Blood smears were taken daily for two weeks after
499 challenge. Parasitaemia was assessed by microscopic examination of Giemsa-stained
500 smears. Data shown is a combination of two independent experiments.

501 **FIG 2** Prime-boost vaccination with viral vectored CSP-expressing vaccines induces strong
502 anti-CSP antibody and CD8+ T cell responses, and SYIPSAEKI-specific CD8+ T cell
503 responses are essential for protection. (A) BALB/c mice were vaccinated with AdCh63 and
504 MVA vaccines expressing *PbCSP* (Ad*PbCSP* and MVA*PbCSP*) and challenged with 10,000

505 *PbWT* or *PbCSP*^{SIINFEKL} sporozoites as shown. (B) Flow cytometry gating strategy used to
 506 determine proportions of IFN- γ + CD11a+ CD8+ T cells. (C) Proportion of IFN- γ -producing
 507 CD11a of total CD8+ T cells. Blood was drawn from the tail from naïve (n=9) and vaccinated
 508 mice (n=10) two weeks after boost and restimulated with SYIPSAEKI and stained for CD8
 509 and CD11a surface markers, and IFN- γ for flow cytometric analysis. (D) Reciprocal antibody
 510 titers of mouse serum reactive to whole sporozoites. Serum from naïve (n=11) and
 511 vaccinated mice (n=12) was isolated two weeks after boost and CSP specific antibody titres
 512 were measured by immunofluorescent antibody assay. (E) Livers from vaccinated mice (+)
 513 challenged with *PbWT* (n=6) or *PbCSP*^{SIINFEKL} sporozoites (n=5) and non-vaccinated mice (-)
 514 challenged with *PbWT* (n=5) or *PbCSP*^{SIINFEKL} sporozoites (n=5) were harvested 42 hours
 515 post-challenge and relative liver parasite levels were quantified using the $\Delta\Delta C_t$ method
 516 comparing levels of *P. berghei* 18S rRNA and levels of mouse *GAPDH* mRNA. (F) Groups of
 517 vaccinated and non-vaccinated mice (n=6) were challenged with 5,000 *PbWT* or
 518 *PbCSP*^{SIINFEKL} sporozoites. Vaccinated mice challenged with *PbWT* (triangles) or
 519 *PbCSP*^{SIINFEKL} (squares) and non-vaccinated mice challenged with *PbWT* (inverted triangles)
 520 or *PbCSP*^{SIINFEKL} (diamonds) had daily tail smears taken from day 3-14 post challenge. Slides
 521 were stained with Giemsa and parasitaemia was assessed by microscopy. (C-E) Each data
 522 point represents one mouse with mean values (\pm SEM) shown and statistics were calculated
 523 using the Mann-Whiney test (*, $p < 0.05$; ***, $p < 0.001$).

524 **FIG 3** Prime-boost vaccination with CSP expressing viruses does not protect C57BL/6 mice,
 525 irrespective of induced antibody titres.

526 (A) C57BL/6 mice were vaccinated with AdCh63 and MVA vaccines *PbCSP* and challenged
 527 with 10,000 *PbWT* or *PbCSP*^{SIINFEKL} sporozoites as shown. (B) Flow cytometry gating
 528 strategy used to determine proportions of IFN- γ + CD11a+ CD8+ T cells. (C) Proportion of
 529 IFN- γ -producing CD11a of total CD8+ T cells. Blood was drawn from the tail from naïve
 530 (n=10) and vaccinated mice (n=10) two weeks after boost was restimulated with SYIPSAEKI
 531 and stained for CD8 and CD11a surface markers, and IFN- γ for flow cytometric analysis. (D)
 532 Reciprocal antibody titres of mouse serum reactive to whole sporozoites. Serum from naïve

533 (n=6) and vaccinated mice (n=9) was isolated two weeks after boost and CSP specific
534 antibody titres were measured by immunofluorescent antibody assay. (E) Livers from groups
535 of 5 mice per condition were harvested 42 hours post-challenge and relative liver parasite
536 levels were quantified using the $\Delta\Delta C_t$ method comparing levels of *P. berghei* 18S rRNA and
537 levels of mouse *GAPDH* mRNA. None of the differences were significant ($p>0.05$). (C-E).
538 Each data point represents one mouse with mean values (\pm SEM) shown and statistics were
539 calculated using the Mann-Whiney test ($***p<0.001$).

540





